

OSTROWSKI INEQUALITIES ON TIME SCALES

MARTIN BOHNER AND THOMAS MATTHEWS

Missouri University of Science and Technology

Department of Mathematics and Statistics

Rolla, MO 65409-0020, USA

EMail: bohner@mst.edu tmnqb@mst.edu

Received: 06 July, 2007

Accepted: 15 February, 2008


Communicated by: S.S. Dragomir

2000 AMS Sub. Class.: 26D15.

Key words: Ostrowski inequality, Montgomery identity, Time scales.

Abstract: We prove Ostrowski inequalities (regular and weighted cases) on time scales and thus unify and extend corresponding continuous and discrete versions from the literature. We also apply our results to the quantum calculus case.

Acknowledgements: The authors thank the referees for their careful reading of the manuscript and insightful comments.





Ostrowski Inequalities on
Time Scales

Martin Bohner and
Thomas Matthews

vol. 9, iss. 1, art. 6, 2008

[Title Page](#)

[Contents](#)



Page 1 of 17

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Contents

1	Introduction	3
2	Time Scales Essentials	4
3	The Ostrowski Inequality on Time Scales	7
4	The Weighted Case	13



Ostrowski Inequalities on
Time Scales

Martin Bohner and
Thomas Matthews

vol. 9, iss. 1, art. 6, 2008

Title Page

Contents



Page 2 of 17


Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



1. Introduction

In 1938, Ostrowski derived a formula to estimate the absolute deviation of a differentiable function from its integral mean. As shown in [6], the so-called Ostrowski inequality

$$(1.1) \quad \left| f(t) - \frac{1}{b-a} \int_a^b f(s) ds \right| \leq \sup_{a < t < b} |f'(t)| (b-a) \left[\frac{\left(t - \frac{a+b}{2}\right)^2}{(b-a)^2} + \frac{1}{4} \right]$$

holds and can be shown by using the Montgomery identity [5]. These two properties will be proved for general time scales, which unify discrete, continuous and many other cases. The setup of this paper is as follows. In Section 2 we first give some preliminary results on time scales that are needed in the remainder of this paper. Next, in Section 3 we prove time scales versions of the Montgomery identity and of the Ostrowski inequality (1.1) (the question of sharpness is also addressed), while in Section 4 we offer several weighted time scales versions of the Ostrowski inequality. Throughout, we apply our results to the special cases of continuous, discrete, and quantum time scales.

Ostrowski Inequalities on
Time Scales
Martin Bohner and
Thomas Matthews
vol. 9, iss. 1, art. 6, 2008

[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 3 of 17

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

2. Time Scales Essentials

Definition 2.1. A time scale is an arbitrary nonempty closed subset of the real numbers.

The most important examples of time scales are \mathbb{R} , \mathbb{Z} and $q^{\mathbb{N}_0} := \{q^k \mid k \in \mathbb{N}_0\}$.

Definition 2.2. If \mathbb{T} is a time scale, then we define the forward jump operator $\sigma : \mathbb{T} \rightarrow \mathbb{T}$ by $\sigma(t) := \inf \{s \in \mathbb{T} \mid s > t\}$ for all $t \in \mathbb{T}$, the backward jump operator $\rho : \mathbb{T} \rightarrow \mathbb{T}$ by $\rho(t) := \sup \{s \in \mathbb{T} \mid s < t\}$ for all $t \in \mathbb{T}$, and the graininess function $\mu : \mathbb{T} \rightarrow [0, \infty)$ by $\mu(t) := \sigma(t) - t$ for all $t \in \mathbb{T}$. Furthermore for a function $f : \mathbb{T} \rightarrow \mathbb{R}$, we define $f^\sigma(t) = f(\sigma(t))$ for all $t \in \mathbb{T}$ and $f^\rho(t) = f(\rho(t))$ for all $t \in \mathbb{T}$. In this definition we use $\inf \emptyset = \sup \mathbb{T}$ (i.e., $\sigma(t) = t$ if t is the maximum of \mathbb{T}) and $\sup \emptyset = \inf \mathbb{T}$ (i.e., $\rho(t) = t$ if t is the minimum of \mathbb{T}).

These definitions allow us to characterize every point in a time scale as displayed in Table 1.

t right-scattered	$t < \sigma(t)$
t right-dense	$t = \sigma(t)$
t left-scattered	$\rho(t) < t$
t left-dense	$\rho(t) = t$
t isolated	$\rho(t) < t < \sigma(t)$
t dense	$\rho(t) = t = \sigma(t)$

Table 1: Classification of Points

Definition 2.3. A function $f : \mathbb{T} \rightarrow \mathbb{R}$ is called rd-continuous (denoted by C_{rd}) if it is continuous at right-dense points of \mathbb{T} and its left-sided limits exist (finite) at left-dense points of \mathbb{T} .



Title Page

Contents

◀◀ ▶▶


◀ ▶

Page 4 of 17

Go Back

Full Screen

Close



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 5 of 17

[Go Back](#)

[Full Screen](#)

[Close](#)

Theorem 2.4 (Existence of Antiderivatives). Let f be rd-continuous. Then f has an antiderivative F satisfying $F^\Delta = f$.

Proof. See [1, Theorem 1.74]. □

Definition 2.5. If f is rd-continuous and $t_0 \in \mathbb{T}$, then we define the integral

$$(2.1) \quad F(t) = \int_{t_0}^t f(\tau) \Delta \tau \quad \text{for } t \in \mathbb{T}.$$

Therefore for $f \in C_{\text{rd}}$ we have $\int_a^b f(\tau) \Delta \tau = F(b) - F(a)$, where $F^\Delta = f$.


Theorem 2.6. Let f, g be rd-continuous, $a, b, c \in \mathbb{T}$ and $\alpha, \beta \in \mathbb{R}$. Then

1. $\int_a^b [\alpha f(t) + \beta g(t)] \Delta t = \alpha \int_a^b f(t) \Delta t + \beta \int_a^b g(t) \Delta t$,
2. $\int_a^b f(t) \Delta t = - \int_b^a f(t) \Delta t$,
3. $\int_a^b f(t) \Delta t = \int_a^c f(t) \Delta t + \int_c^b f(t) \Delta t$,
4. $\int_a^b f(t) g^\Delta(t) \Delta t = (fg)(b) - (fg)(a) - \int_a^b f^\Delta(t) g(\sigma(t)) \Delta t$,
5. $\int_a^a f(t) \Delta t = 0$.

Proof. See [1, Theorem 1.77]. □

Definition 2.7. Let $g_k, h_k : \mathbb{T}^2 \rightarrow \mathbb{R}$, $k \in \mathbb{N}_0$ be defined by

$$g_0(t, s) = h_0(t, s) = 1 \quad \text{for all } s, t \in \mathbb{T}$$



Title Page

Contents



Page 6 of 17

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

and then recursively by

$$g_{k+1}(t, s) = \int_s^t g_k(\sigma(\tau), s) \Delta\tau \quad \text{for all } s, t \in \mathbb{T}$$

and


$$h_{k+1}(t, s) = \int_s^t h_k(\tau, s) \Delta\tau \quad \text{for all } s, t \in \mathbb{T}.$$

Theorem 2.8 (Hölder's Inequality). Let $a, b \in \mathbb{T}$ and $f, g : [a, b] \rightarrow \mathbb{R}$ be rd-continuous. Then

$$(2.2) \quad \int_a^b |f(t)g(t)| \Delta t \leq \left\{ \int_a^b |f(t)|^p \Delta t \right\}^{\frac{1}{p}} \left\{ \int_a^b |g(t)|^q \Delta t \right\}^{\frac{1}{q}},$$

where $p > 1$ and $\frac{1}{p} + \frac{1}{q} = 1$.

Proof. See [1, Theorem 6.13]. □



3. The Ostrowski Inequality on Time Scales

Lemma 3.1 (Montgomery Identity). Let $a, b, s, t \in \mathbb{T}$, $a < b$ and $f : [a, b] \rightarrow \mathbb{R}$ be differentiable. Then

$$(3.1) \quad f(t) = \frac{1}{b-a} \int_a^b f^\sigma(s) \Delta s + \frac{1}{b-a} \int_a^b p(t, s) f^\Delta(s) \Delta s,$$

where

$$(3.2) \quad p(t, s) = \begin{cases} s - a, & a \leq s < t, \\ s - b, & t \leq s \leq b. \end{cases}$$

Proof. Using Theorem 2.6 (4), we have

$$\int_a^t (s - a) f^\Delta(s) \Delta s = (t - a) f(t) - \int_a^t f^\sigma(s) \Delta s$$

and similarly

$$\int_t^b (s - b) f^\Delta(s) \Delta s = (b - t) f(t) - \int_t^b f^\sigma(s) \Delta s.$$

Therefore

$$\begin{aligned} & \frac{1}{b-a} \int_a^b f^\sigma(s) \Delta s + \frac{1}{b-a} \int_a^b p(t, s) f^\Delta(s) \Delta s \\ &= \frac{1}{b-a} \int_a^b f^\sigma(s) \Delta s + \frac{1}{b-a} \left[(b-a) f(t) - \int_a^b f^\sigma(s) \Delta s \right] \\ &= f(t), \end{aligned}$$

i.e., (3.1) holds. □

[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 7 of 17


[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 8 of 17

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.

If we apply Lemma 3.1 to the discrete and continuous cases, we have the following results.

Corollary 3.2 (discrete case). *We let $\mathbb{T} = \mathbb{Z}$. Let $a = 0$, $b = n$, $s = j$, $t = i$ and $f(k) = x_k$. Then*

$$x_i = \frac{1}{n} \sum_{j=1}^n x_j + \frac{1}{n} \sum_{j=0}^{n-1} p(i, j) \Delta x_j,$$

where


$$\begin{aligned} p(i, 0) &= 0, \\ p(1, j) &= j - n \quad \text{for } 1 \leq j \leq n - 1, \\ p(n, j) &= j \quad \text{for } 0 \leq j \leq n - 1, \\ p(i, j) &= \begin{cases} j, & 0 \leq j < i, \\ j - n, & i \leq j \leq n - 1 \end{cases} \end{aligned}$$

as we just need $1 \leq i \leq n$ and $0 \leq j \leq n - 1$. This result is the same as in [2, Theorem 2.1].

Corollary 3.3 (continuous case). *We let $\mathbb{T} = \mathbb{R}$. Then*



$$f(t) = \frac{1}{b-a} \int_a^b f(s) ds + \frac{1}{b-a} \int_a^b p(t, s) f'(s) ds.$$

This is the Montgomery identity in the continuous case, which can be found in [5, p. 565].



Title Page

Contents



Page 9 of 17

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Corollary 3.4 (quantum calculus case). We let $\mathbb{T} = q^{\mathbb{N}_0}$, $q > 1$, $a = q^m$ and $b = q^n$ with $m < n$. Then

$$f(t) = \frac{\sum_{k=m}^{n-1} q^k f(q^{k+1})}{\sum_{k=m}^{n-1} q^k} + \frac{1}{q^n - q^m} \sum_{k=m}^{n-1} [f(q^{k+1}) - f(q^k)] p(t, q^k),$$

where

$$p(t, q^k) = \begin{cases} q^k - q^m, & q^m \leq q^k < t, \\ q^k - q^n, & t \leq q^k \leq q^n. \end{cases}$$

Theorem 3.5 (Ostrowski Inequality). Let $a, b, s, t \in \mathbb{T}$, $a < b$ and $f : [a, b] \rightarrow \mathbb{R}$ be differentiable. Then

$$(3.3) \quad \left| f(t) - \frac{1}{b-a} \int_a^b f^\sigma(s) \Delta s \right| \leq \frac{M}{b-a} (h_2(t, a) + h_2(t, b)),$$

where

$$M = \sup_{a < t < b} |f^\Delta(t)|.$$

This inequality is sharp in the sense that the right-hand side of (3.3) cannot be replaced by a smaller one.

Proof. With Lemma 3.1 and $p(t, s)$ defined as in (3.2), we have

$$\begin{aligned} \left| f(t) - \frac{1}{b-a} \int_a^b f^\sigma(s) \Delta s \right| &= \left| \frac{1}{b-a} \int_a^b p(t, s) f^\Delta(s) \Delta s \right| \\ &\leq \frac{M}{b-a} \left(\int_a^t |s-a| \Delta s + \int_t^b |s-b| \Delta s \right) \end{aligned}$$

$$\begin{aligned}
&= \frac{M}{b-a} \left(\int_a^t (s-a) \Delta s + \int_t^b (b-s) \Delta s \right) \\
&= \frac{M}{b-a} (h_2(t, a) + h_2(t, b)).
\end{aligned}$$

Note that, since $p(t, a) = 0$, the smallest value attaining the supremum in M is greater than a . To prove the sharpness of this inequality, let $f(t) = t$, $a = t_1$, $b = t_2$ and $t = t_2$. It follows that $f^\Delta(t) = 1$ and $M = 1$. Starting with the left-hand side of (3.3), we have

$$\begin{aligned}
\left| f(t) - \frac{1}{b-a} \int_a^b f^\sigma(s) \Delta s \right| &= \left| t_2 - \frac{1}{t_2 - t_1} \int_{t_1}^{t_2} \sigma(s) \Delta s \right| \\
&= \left| t_2 - \frac{1}{t_2 - t_1} \left(\int_{t_1}^{t_2} (\sigma(s) + s) \Delta s - \int_{t_1}^{t_2} s \Delta s \right) \right| \\
&= \left| t_2 - \frac{1}{t_2 - t_1} \left(\int_{t_1}^{t_2} (s^2)^\Delta \Delta s - \int_{t_1}^{t_2} s \Delta s \right) \right| \\
&= \left| -t_1 + \frac{1}{t_2 - t_1} \int_{t_1}^{t_2} s \Delta s \right|.
\end{aligned}$$

Starting with the right-hand side of (3.3), we have

$$\begin{aligned}
\frac{M}{b-a} (h_2(t, a) + h_2(t, b)) &= \frac{1}{t_2 - t_1} \left(\int_{t_1}^{t_2} (s - t_1) \Delta s + \int_{t_2}^{t_2} (s - t_2) \Delta s \right) \\
&= \frac{1}{t_2 - t_1} \left(-t_1 t_2 + t_1^2 + \int_{t_1}^{t_2} s \Delta s \right) \\
&= -t_1 + \frac{1}{t_2 - t_1} \int_{t_1}^{t_2} s \Delta s.
\end{aligned}$$



Title Page

Contents

◀ ▶

◀ ▶

Page 10 of 17


Go Back

Full Screen

Close



journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



[Title Page](#)

[Contents](#)



Page 11 of 17

[Go Back](#)

[Full Screen](#)

[Close](#)

Therefore in this particular case

$$\left| f(t) - \frac{1}{b-a} \int_a^b f^\sigma(s) \Delta s \right| \geq \frac{M}{b-a} (h_2(t, a) + h_2(t, b))$$

and by (3.3) also

$$\left| f(t) - \frac{1}{b-a} \int_a^b f^\sigma(s) \Delta s \right| \leq \frac{M}{b-a} (h_2(t, a) + h_2(t, b)).$$

So the sharpness of the Ostrowski inequality is shown. \square

Corollary 3.6 (discrete case). Let $\mathbb{T} = \mathbb{Z}$. Let $a = 0$, $b = n$, $s = j$, $t = i$ and $f(k) = x_k$. Then

$$(3.4) \quad \left| x_i - \frac{1}{n} \sum_{j=1}^n x_j \right| \leq \frac{M}{n} \left[\left| i - \frac{n+1}{2} \right|^2 + \frac{n^2-1}{4} \right],$$


where

$$M = \max_{1 \leq i \leq n-1} |\Delta x_i|.$$

This is the discrete Ostrowski inequality from [2, Theorem 3.1], where the constant $\frac{1}{4}$ in the right-hand side of (3.4) is the best possible in the sense that it cannot be replaced by a smaller one.

Corollary 3.7 (continuous case). If $\mathbb{T} = \mathbb{R}$, then

$$\left| f(t) - \frac{1}{b-a} \int_a^b f(s) ds \right| \leq M(b-a) \left[\frac{(t - \frac{a+b}{2})^2}{(b-a)^2} + \frac{1}{4} \right],$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 12 of 17

Go Back

Full Screen

Close

where

$$M = \sup_{a < t < b} |f'(t)|.$$

This is the Ostrowski inequality in the continuous case [6, p. 226–227], where again the constant $\frac{1}{4}$ in the right-hand side is the best possible.


Corollary 3.8 (quantum calculus case). Let $\mathbb{T} = q^{\mathbb{N}_0}$, $q > 1$, $a = q^m$ and $b = q^n$ with $m < n$. Then

$$\left| f(t) - \frac{1}{q^n - q^m} \int_{q^m}^{q^n} f^\sigma(s) \Delta s \right| \leq \frac{M}{q^n - q^m} \left[\frac{2}{1+q} \left(\left(t - \frac{\frac{1+q}{2}(q^m + q^n)}{2} \right)^2 + \frac{(\frac{1+q}{2})^2 (q^m + q^n)^2 + (2(1+q) - 2)(q^{2m} + q^{2n})}{4} \right) \right],$$

where

$$M = \sup_{q^m < t < q^n} \left| \frac{f(qt) - f(t)}{(q-1)t} \right|,$$

and the constant $\frac{1}{4}$ in the right-hand side is the best possible.



Title Page

Contents



Page 13 of 17

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

4. The Weighted Case

The following weighted Ostrowski inequality on time scales holds.

Theorem 4.1. Let $a, b, s, t, \tau \in \mathbb{T}$, $a < b$ and $f : [a, b] \rightarrow \mathbb{R}$ be differentiable, $q \in C_{\text{rd}}$. Then

$$(4.1) \quad \left| f(t) - \int_a^b q^\sigma(s) f^\sigma(s) \Delta s \right| \leq \int_a^b q^\sigma(s) |\sigma(s) - t| M \Delta s$$

$$(4.2) \quad \leq M \begin{cases} \left(\int_a^b |\sigma(s) - t|^p \Delta s \right)^{\frac{1}{p}} \left(\int_a^b (q^\sigma(s))^q \Delta s \right)^{\frac{1}{q}}, & \frac{1}{p} + \frac{1}{q} = 1, \quad p > 1; \\ \sup_{a \leq s < b} q^\sigma(s) [g_2(a, t) + g_2(b, t)]; \\ \frac{b - \sigma(a)}{2} + \left| t - \frac{b + \sigma(a)}{2} \right|, \end{cases}$$

where

$$M = \sup_{\sigma(a) \leq \tau < b} |f^\Delta(\tau)|$$

and

$$\int_a^b q^\sigma(s) \Delta s = 1, \quad q(s) \geq 0.$$

Proof. We have

$$\begin{aligned} \left| f(t) - \int_a^b q^\sigma(s) f^\sigma(s) \Delta s \right| &= \left| \int_a^b q^\sigma(s) (f(t) - f^\sigma(s)) \Delta s \right| \end{aligned}$$



[Title Page](#)

[Contents](#)

◀◀ ▶▶

◀ ▶

Page 14 of 17

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.

$$\begin{aligned} &\leq \int_a^t q^\sigma(s) |f(t) - f^\sigma(s)| \Delta s + \int_t^b q^\sigma(s) |f(t) - f^\sigma(s)| \Delta s \\ &\leq \int_a^t q^\sigma(s) \int_{\sigma(s)}^t |f^\Delta(\tau)| \Delta \tau \Delta s + \int_t^b q^\sigma(s) \int_t^{\sigma(s)} |f^\Delta(\tau)| \Delta \tau \Delta s \\ &\leq M \int_a^b q^\sigma(s) |\sigma(s) - t| \Delta s, \end{aligned}$$

and therefore (4.1) is shown.

The first part of (4.2) can be done easily by applying (2.2). By factoring $\sup_{a \leq s < b} q^\sigma(s)$, we have


$$\begin{aligned} \int_a^b q^\sigma(s) |\sigma(s) - t| \Delta s &\leq \sup_{a \leq s < b} q^\sigma(s) \left(\int_a^t (t - \sigma(s)) \Delta s + \int_t^b (\sigma(s) - t) \Delta s \right) \\ &= \sup_{a \leq s < b} q^\sigma(s) \left(\int_t^a (\sigma(s) - t) \Delta s + \int_t^b (\sigma(s) - t) \Delta s \right) \\ &= \sup_{a \leq s < b} q^\sigma(s) [g_2(a, t) + g_2(b, t)], \end{aligned}$$

and therefore the second part of (4.2) holds. Finally for proving the third inequality, we use the fact that

$$\sup_{a \leq s < b} \{|\sigma(s) - t|\} = \max \{b - t, t - \sigma(a)\} = \frac{b - \sigma(a)}{2} + \left| t - \frac{b + \sigma(a)}{2} \right|.$$



Thus (4.2) is shown. □

Remark 1. Theorem 4.1 states a similar result as shown in [3, Theorem 3.1], if we consider the normalized isotonic functional $A(f) = \int_a^b q^\sigma(s) f^\sigma(s) \Delta s$. Moreover



[Title Page](#)

[Contents](#)



Page 15 of 17

[Go Back](#)

[Full Screen](#)

[Close](#)

the second inequality of (4.2) is comparable to the achievement in [4, Theorem 3.1] for the continuous case (see [4, Corollary 3.3]).

Corollary 4.2 (discrete case). Let $\mathbb{T} = \mathbb{Z}$. Let $a = 0$, $b = n$, $s = j$, $t = i$, $\tau = k$ and $f(k) = x_k$. Then $\sum_{i=1}^n q_i = 1$, $q_i \leq 0$ and

$$\left| x_i - \sum_{j=1}^n q_j x_j \right| \leq \sum_{j=1}^n q_j |j - i| M$$

$$\leq M \begin{cases} \left(\sum_{j=1}^n |j - i|^p \right)^{\frac{1}{p}} \left(\sum_{j=1}^n q_j^q \right)^{\frac{1}{q}}, & \frac{1}{p} + \frac{1}{q} = 1, \quad p > 1; \\ \left[\frac{n^2-1}{4} + \left(i - \frac{n+1}{2} \right)^2 \right] \max_{j=1..n} q(j); \\ \frac{n-1}{2} + \left| i - \frac{n+1}{2} \right|, \end{cases}$$

where


$$M = \max_{k=1..n-1} |\Delta x_k|.$$

This is the result given in [2, Theorem 4.1].

Corollary 4.3 (continuous case). If $\mathbb{T} = \mathbb{R}$, then $\int_a^b q(s) ds = 1$, $q(s) \geq 0$ and

$$\left| f(t) - \int_a^b q(s) f(s) ds \right| \leq \int_a^b q(s) |s - t| M ds$$

$$\leq M \begin{cases} \left(\int_a^b |s - t|^p ds \right)^{\frac{1}{p}} \left(\int_a^b (q(s))^q ds \right)^{\frac{1}{q}}, & \frac{1}{p} + \frac{1}{q} = 1, \quad p > 1; \\ \sup_{a \leq s < b} q(s) (b - a)^2 \left[\frac{\left(t - \frac{a+b}{2} \right)^2}{(b-a)^2} + \frac{1}{4} \right]; \\ \frac{b-a}{2} + \left| t - \frac{b+a}{2} \right|, \end{cases}$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 16 of 17

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

© 2007 Victoria University. All rights reserved.

where

$$M = \sup_{a \leq \tau < b} |f'(\tau)|.$$

Another interesting conclusion of Theorem 4.1 is the following corollary.

Corollary 4.4. *Let $a, b, s, t \in \mathbb{T}$ and f differentiable. Then*

$$(4.3) \quad \left| f(t) - \frac{1}{b-a} \int_a^b f^\sigma(s) \Delta s \right| \leq \frac{M}{b-a} (h_2(t, a) + h_2(t, b)),$$

where

$$M = \sup_{\sigma(a) \leq t < b} |f^\Delta(t)|.$$

Note that this was shown in a different manner in Theorem 3.5. In (4.3) we use the fact that the functions g_2 and h_2 satisfy $g_2(s, t) = (-1)^2 h_2(t, s)$ for all $t \in \mathbb{T}$ and all $s \in \mathbb{T}^\kappa$ (see [1, Theorem 1.112]).

Remark 2. Moreover note that there is a small difference of (4.2) in comparison to Theorem 3.5, as we have $\sup_{\sigma(a) \leq t < b}$ instead of $\sup_{a < t < b}$. This is just important if a is right-dense, i.e., $\sigma(a) = a$. But in those cases the inequality does not change and is still sharp. Furthermore in the proof of Theorem 3.5 we could have picked $\sup_{a \leq t < b}$ as explained before.

References

- [1] M. BOHNER AND A. PETERSON, *Dynamic equations on time scales*, Boston, MA: Birkhäuser Boston Inc., 2001.
- [2] S.S. DRAGOMIR, The discrete version of Ostrowski's inequality in normed linear spaces, *J. Inequal. Pure Appl. Math.*, **3**(1) (2002), Art. 2. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=155>]
- [3] S.S. DRAGOMIR, Ostrowski type inequalities for isotonic linear functionals, *J. Inequal. Pure Appl. Math.*, **3**(5) (2002), Art. 68. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=220>]
- [4] B. GAVREA AND I. GAVREA, Ostrowski type inequalities from a linear functional point of view, *J. Inequal. Pure Appl. Math.*, **1**(2) (2000), Art. 11. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=104>]
- [5] D.S. MITRINOVIĆ, J.E. PEČARIĆ AND A.M. FINK, *Inequalities involving functions and their integrals and derivatives*, vol. 53 of *Mathematics and its Applications (East European Series)*. Dordrecht: Kluwer Academic Publisher Group, p. 565, 1991.
- [6] A. OSTROWSKI, Über die Absolutabweichung einer differenzierbaren Funktion von ihrem Integralmittelwert, *Comment. Math. Helv.*, **10**(1) (1937), 226–227.




Ostrowski Inequalities on
Time Scales

Martin Bohner and
Thomas Matthews

vol. 9, iss. 1, art. 6, 2008

Title Page

Contents



Page 17 of 17

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756