

## GEOMETRY AND TOPOLOGY OF COADJOINT ORBITS OF SEMISIMPLE LIE GROUPS

JULIA BERNATSKA and PETRO HOLOD

*Department of Physical and Mathematical Sciences, National University of  
Kyiv-Mohyla Academy, 2 Skovorody Str., 04070 Kyiv, Ukraine*

**Abstract.** Orbits of coadjoint representations of classical compact Lie groups have a lot of applications. They appear in representation theory, geometrical quantization, theory of magnetism, quantum optics, etc. As geometric objects the orbits were the subject of extensive study. However, they remain hard for calculation and application. We propose a simple solution for the following problem: an explicit parametrization of the orbit by means of a generalized stereographic projection, which provide a Kählerian structure on the orbit, and basis two-forms for the cohomology group of the orbit.

### 1. Introduction

Orbits of coadjoint representations of semisimple Lie groups are an extremely interesting subject. These homogeneous spaces are flag manifolds. Remarkable, that the coadjoint orbits of compact groups are Kählerian manifolds. In 1950s Borel, Bott, Koszul, Hirzebruch et al. investigated the coadjoint orbits as complex homogeneous manifolds. It was proven that each coadjoint orbit of a compact connected Lie group  $G$  admits a canonical  $G$ -invariant complex structure and the only (within homotopies)  $G$ -invariant Kählerian metrics. Furthermore, the coadjoint orbits can be considered as fibre bundles whose bases and fibres are coadjoint orbits themselves.

Coadjoint orbits appear in many spheres of theoretical physics, for instance in representation theory, geometrical quantization, theory of magnetism, quantum optics. They serve as definitional domains in problems connected with nonlinear integrable equations (so called equations of soliton type). Since these equations have a wide application, the remarkable properties of coadjoint orbits interest not only mathematicians but also physicists.

It should be pointed out that much of our material is, of course, not new, but drawn from various areas of the mathematical literature. The material was collected for solving the physical problem based on a classical Heisenberg equation with  $SU(n)$  as a gauge group. The equation describes a behavior of magnetics with spin  $s \geq 1$ . The paper includes an investigation of geometrical and topological properties of the coadjoint orbits. We hope it fulfills a certain need. We would like to mention that we have added a number of new results (such as an explicit expression for a stereographic projection in the case of group  $SU(3)$  and improving the way of its computation, the idea of obtaining the Kählerian potential on an orbit, an introduction of basis two-forms for the cohomology ring of an orbit).

The paper is organized as follows. In Section 2 we recall the notion of a coadjoint orbit, propose a classification of the orbits, and describe the orbit as a fibre bundle over an orbit with an orbit as a fibre. Section 3 is devoted to a generalized stereographic projection from a Lie algebra onto its coadjoint orbit, it gives a suitable complex parametrization of the orbit. As an example, we compute an explicit expression for the stereographic projection in the case of group  $SU(3)$ . In Section 4 we propose a way of obtaining Kählerian structures and Kählerian potentials on the orbits. Section 5 concerns a structure of the cohomology rings of the orbits and finding of  $G$ -invariant bases for the cohomology groups.

## 2. Coadjoint Orbits of Semisimple Lie Groups

We start with recalling the notion of a coadjoint orbit. Let  $G$  be a compact semisimple classical Lie group,  $\mathfrak{g}$  denote the corresponding Lie algebra, and  $\mathfrak{g}^*$  denote the dual space of  $\mathfrak{g}$ . Let  $T$  be the maximal torus of  $G$ , and  $\mathfrak{h}$  be the maximal commutative subalgebra (also called a Cartan subalgebra) of  $\mathfrak{g}$ . Accordingly,  $\mathfrak{h}^*$  denotes the dual space of  $\mathfrak{h}$ .

**Definition 1.** *The subset  $\mathcal{O}_\mu = \{Ad_g^* \mu; g \in G\}$  of  $\mathfrak{g}^*$  is called a **coadjoint orbit** of  $G$  through  $\mu \in \mathfrak{g}^*$ .*

In the case of classical Lie groups we can use the standard representations for adjoint and coadjoint operators

$$Ad_g X = gXg^{-1}, \quad X \in \mathfrak{g}, \quad Ad_g^* \mu = g^{-1} \mu g, \quad \mu \in \mathfrak{g}^*.$$

Comparing these formulas one can easily see that *a coadjoint orbit coincides with the adjoint.*

Define the *stability subgroup* at a point  $\mu \in \mathfrak{g}^*$  as  $G_\mu = \{g \in G; Ad_g^* \mu = \mu\}$ . The coadjoint operator induces a bijective correspondence between an orbit  $\mathcal{O}_\mu$  and a coset space  $G_\mu \backslash G$  (in the sequel, we deal with right coset spaces).

First of all, we classify the coadjoint orbits of an arbitrary semisimple group  $G$ . Obviously, each orbit is drawn from a unique point, which we call an *initial point*

and denote by  $\mu_0$ . The following theorem from [1] allows to restrict the region of search of an initial point.

**Theorem ([1]).** *Each orbit of the coadjoint action of  $G$  intersects  $\mathfrak{h}^*$  precisely in an orbit of the Weyl group.*

In other words, each orbit is assigned to a finite non-empty subset of  $\mathfrak{h}^*$ . For more detail recall the notion of the Weyl group. Let  $N(H)$  be the **normalizer** of a subset  $H \subset G$  in  $G$ , that is  $N(H) = \{g \in H; g^{-1}Hg = H\}$ . Let  $C(H)$  be the **centralizer** of  $H$ , that is  $C(H) = \{g \in G; g^{-1}hg = h, h \in H\}$ . Obviously,  $C(T) = T$ , where  $T$  is the maximal torus of  $G$ .

**Definition 2.** *The **Weyl group** of  $G$  is the factor-group of  $N(T)$  over  $C(T)$*

$$W(G) = N(T)/C(T).$$

The Weyl group  $W(G)$  acts transitively on  $\mathfrak{h}^*$ . The action of  $W(G)$  is performed by the coadjoint operator. It is easy to show that  $W(G)$  is isomorphic to the finite group generated by reflections  $w_\alpha$  across the hyperplanes orthogonal to simple roots  $\alpha$

$$w_\alpha(\mu) = \mu - 2 \frac{\langle \mu, \alpha \rangle}{\langle \alpha, \alpha \rangle} \alpha, \quad \mu \in \mathfrak{h}^*$$

where  $\langle \cdot, \cdot \rangle$  denotes a bilinear form on  $\mathfrak{g}^*$ .

**Definition 3.** *The open domain*

$$C = \{\mu \in \mathfrak{h}^*; \langle \mu, \alpha \rangle > 0, \alpha \in \Delta^+\}$$


is called the **positive Weyl chamber**. Here  $\Delta^+$  denotes the set of positive roots.

We call the set  $\Gamma_\alpha = \{\mu \in \mathfrak{h}^*; \langle \mu, \alpha \rangle = 0\}$  a **wall of the Weyl chamber**.

If we reflect the closure  $\overline{C}$  of the positive Weyl chamber by elements of the Weyl group we cover  $\mathfrak{h}^*$  overall

$$\mathfrak{h}^* = \bigcup_{w \in W(G)} w \cdot \overline{C}.$$

An orbit of the Weyl group  $W(G)$  is obtained by the action of  $W(G)$  on a point of  $\overline{C}$ . In the case of group  $SU(3)$ , two possible types of orbits of the Weyl group are shown on the root diagram (see Fig. 1). Black points denote intersections of a coadjoint orbit with  $\mathfrak{h}^*$  and form an orbit of  $W(SU(3))$ . The positive Weyl chamber is filled with grey color. It has two walls:  $\Gamma_{\alpha_1}$  and  $\Gamma_{\alpha_2}$ . The respective reflections across these hyperplanes are denoted by  $w_{\alpha_1}$  and  $w_{\alpha_2}$ . At the left, one can see a generic case, when an orbit of  $W(SU(3))$  has six elements. It happens if an initial point lies in the interior of the positive Weyl chamber. At the right, there is a degenerate (non-generic) case, when an orbit of  $W(SU(3))$  has three elements. It happens if an initial point belongs to a wall of the positive Weyl chamber.



**Figure 1.** Root diagram for  $SU(3)$ .

In the both cases the closed positive Weyl chamber contains a unique point of an orbit of  $W(G)$ . We obtain the following

**Proposition 1.** *Each orbit  $\mathcal{O}$  of  $G$  is uniquely defined by an initial point  $\mu_0 \in \mathfrak{h}^*$ , which is located in the closed positive Weyl chamber  $\bar{C}$ . If  $\mu_0$  lies in the interior of the positive Weyl chamber:  $\mu_0 \in C$ , it gives rise to a **generic orbit**. If  $\mu_0$  belongs to a wall of the positive Weyl chamber:  $\mu_0 \in \Gamma_\alpha$ ,  $\alpha \in \Delta^+$ , it gives rise to a **degenerate orbit**.*

As mentioned above, one can define the orbit  $\mathcal{O}_{\mu_0}$  through an initial point  $\mu_0 \in \mathfrak{h}^*$  by  $\mathcal{O}_{\mu_0} = G_{\mu_0} \backslash G$ . Note, that a stability subgroup  $G_\mu$  as  $\mu \in \mathfrak{h}^*$  generically coincides with the maximal torus  $T$ . However, if  $\mu$  belongs to a degenerate orbit, then  $G_\mu$  is a larger subgroup of  $G$  containing  $T$ . Therefore, we define a generic orbit by

$$\mathcal{O}_{\mu_0} = T \backslash G$$

and a degenerate one by

$$\mathcal{O}_{\mu_0} = G_{\mu_0} \backslash G$$

where  $G_{\mu_0} \neq T$ ,  $G_{\mu_0} \supset T$ .

An important topological property of the coadjoint orbits is the following. *Almost each orbit can be regarded as a fibre bundle over an orbit with an orbit as a fibre, except for the maximal degenerate orbits.* Indeed, if there exists an initial point  $\mu_0$  such that  $G_{\mu_0} \supset T$ , one can form a coset space  $T \backslash G_{\mu_0}$ . Thus, the orbit  $\mathcal{O}_{\mu_0} = T \backslash G$  is a fibre bundle over the base  $G_{\mu_0} \backslash G$  with the fibre  $T \backslash G_{\mu_0}$

$$\mathcal{O}_{\mu_0} = \mathcal{E}(G_{\mu_0} \backslash G, T \backslash G_{\mu_0}, \pi)$$

where  $\pi$  denotes a projection from the orbit onto the base. Moreover,  $G_{\mu_0} \backslash G$  and  $T \backslash G_{\mu_0}$  are coadjoint orbits themselves. We formulate this as

**Proposition 2.** *Suppose  $\mathcal{O}_{\mu_0} = G_{\mu_0} \backslash G$  is not the maximal degenerate orbit of  $G$ . Then a subgroup  $K$  such that  $G \supset K \supset G_{\mu_0}$  exists, and  $\mathcal{O}_{\mu_0}$  is a fibre bundle over the base  $K \backslash G$  with the fibre  $G_{\mu_0} \backslash K$*

$$\mathcal{O}_{\mu_0} = \mathcal{E}(K \backslash G, G_{\mu_0} \backslash K, \pi).$$

We will illustrate the proposition by some examples.

**Example 1.** The group  $SU(2)$  has the only type of orbits:

$$\mathcal{O}^{SU(2)} = \frac{SU(2)}{U(1)} \simeq \mathbb{C}P^1.$$

The group  $SU(3)$  has generic and degenerate orbits

$$\mathcal{O}^{SU(3)} = \frac{SU(3)}{U(1) \times U(1)}, \quad \mathcal{O}_d^{SU(3)} = \frac{SU(3)}{SU(2) \times U(1)} \simeq \mathbb{C}P^2.$$

Comparing the above coset spaces we see that a generic orbit  $\mathcal{O}^{SU(3)}$  is a fibre bundle over a degenerate orbit  $\mathcal{O}_d^{SU(3)}$  with a fibre  $\mathcal{O}^{SU(2)}$

$$\mathcal{O}^{SU(3)} = \mathcal{E}(\mathcal{O}_d^{SU(3)}, \mathcal{O}^{SU(2)}, \pi) = \mathcal{E}(\mathbb{C}P^2, \mathbb{C}P^1, \pi).$$

The group  $SU(4)$  has several types of degenerate orbits. There is a list of all possible types of orbits

$$\begin{aligned} \mathcal{O}^{SU(4)} &= \frac{SU(4)}{U(1) \times U(1) \times U(1)}, & \mathcal{O}_{d1}^{SU(4)} &= \frac{SU(4)}{SU(2) \times U(1) \times U(1)} \\ \mathcal{O}_{d2}^{SU(4)} &= \frac{SU(4)}{S(U(2) \times U(2))}, & \mathcal{O}_{d3}^{SU(4)} &= \frac{SU(4)}{SU(3) \times U(1)} \simeq \mathbb{C}P^3. \end{aligned}$$

As a result, there exist several representations of a generic orbit  $\mathcal{O}^{SU(4)}$  as a fibre bundle. For example,

$$\begin{aligned} \mathcal{O}^{SU(4)} &= \mathcal{E}(\mathcal{O}_{d3}^{SU(4)}, \mathcal{O}^{SU(3)}, \pi) = \mathcal{E}(\mathbb{C}P^3, \mathcal{O}^{SU(3)}, \pi) \\ \mathcal{O}^{SU(4)} &= \mathcal{E}(\mathcal{O}_{d2}^{SU(4)}, \mathcal{O}^{SU(2)}, \pi) = \mathcal{E}(\mathcal{O}_{d2}^{SU(4)}, \mathbb{C}P^1, \pi). \end{aligned}$$

**Example 2.** In this paper we consider compact classical Lie groups. They describe linear transformations of real, complex, and quaternionic spaces. Respectively, these groups are  $SO(n)$  over the real field,  $SU(n)$  over the complex field, and  $Sp(n)$  over the quaternionic ring. Here we list the maximal tori of all these groups, and their representations as fibre bundles.

The maximal torus of  $SU(n)$  is  $T = \overbrace{U(1) \times U(1) \times \cdots \times U(1)}^{n-1}$  and the generic type of orbits can be represented as

$$\mathcal{O}^{SU(n)} = \mathcal{E}(\mathbb{C}P^{n-1}, \mathcal{O}^{SU(n-1)}, \pi).$$

The maximal torus of  $SO(n)$  as  $n = 2m$  and  $n = 2m + 1$  has the following form  $T = \underbrace{SO(2) \times SO(2) \times \cdots \times SO(2)}_m$  and in this case the generic type of orbits can be represented as

$$\begin{aligned} \mathcal{O}^{SO(2m)} &= \mathcal{E} \left( G_{2m;2}, \mathcal{O}^{SO(2m-2)}, \pi \right) \\ \mathcal{O}^{SO(2m+1)} &= \mathcal{E} \left( G_{2m-1;2}, \mathcal{O}^{SO(2m-1)}, \pi \right) \end{aligned}$$

where  $G_{2m;2}, G_{2m-1;2}$  denote real Grassman manifolds.

The maximal torus of  $Sp(n)$  is  $T = \overbrace{U(1) \times U(1) \times \cdots \times U(1)}^{n-1}$  while the generic type of orbits can be represented as

$$\mathcal{O}^{Sp(n)} = \mathcal{E} \left( \mathbb{H}P^{n-1}, \mathcal{O}^{Sp(n-1)}, \pi \right)$$

where  $\mathbb{H}$  denotes the quaternionic ring.

### 3. Complex Parameterization of Coadjoint Orbits

In the theory of Lie groups and Lie algebras different ways of parameterization of coadjoint orbits are available. As the most prevalent we choose a **generalized stereographic projection** [2]. It is named so since in the case of group  $SU(2)$  it gives the well-known stereographic projection onto the complex plane, which is the only orbit of  $SU(2)$ . The generalized stereographic projection is a projection from a dual space onto a coadjoint orbit parameterized by complex coordinates.

Complex coordinates are introduced by the well-known procedure that combines Iwasawa and Gauss-Bruhat decompositions. These coordinates are often called **Bruhat coordinates** [3].

We start with complexifying a group  $G$  in the usual way:  $G^{\mathbb{C}} = \exp\{\mathfrak{g} + i\mathfrak{g}\}$ . A generic orbit of  $G$  is defined in  $G^{\mathbb{C}}$  by Montgomery's diffeomorphism

$$\mathcal{O} = T \backslash G \simeq P \backslash G^{\mathbb{C}} \tag{1}$$

where  $P$  denotes the minimal parabolic subgroup of  $G^{\mathbb{C}}$ .

Equation (1) becomes apparent from the Iwasawa decomposition  $G^{\mathbb{C}} = NAK$ , where  $A \simeq \exp\{i\mathfrak{h}\}$  is the real abelian subgroup of  $G^{\mathbb{C}}$ ,  $N$  is a nilpotent subgroup of  $G^{\mathbb{C}}$ , and  $K$  is the maximal compact subgroup of  $G^{\mathbb{C}}$ . Since we consider only compact groups  $G$ ,  $K$  coincides with  $G$ . Then the Iwasawa decomposition of  $G^{\mathbb{C}}$  has the following form

$$G^{\mathbb{C}} = NAG.$$

It is easy to express  $A$  and  $N$  in terms of root vectors. Let  $\Delta^+$  be the set of positive roots  $\alpha$  of  $G^{\mathbb{C}}$ . By  $X_{\alpha}, X_{-\alpha}, \alpha \in \Delta^+$ , denote positive and negative root vectors, respectively. By  $H_{\alpha}, \alpha \in \Delta^+$ , denote the corresponding Cartan vectors, which

form a basis for the Cartan subalgebra  $\mathfrak{h}$ . According to [4], we choose  $X_\alpha$  and  $X_{-\alpha}$  so that  $X_\alpha - X_{-\alpha}, i(X_\alpha + X_{-\alpha}) \in \mathfrak{g}$ . Then

$$N \simeq \exp \left\{ \sum_{\alpha \in \Delta^+} n_\alpha X_\alpha \right\}, \quad n_\alpha \in \mathbb{C}, \quad A \simeq \exp \left\{ \sum_{\alpha \in \Delta^+} a_\alpha iH_\alpha \right\}, \quad a_\alpha \in \mathbb{R}.$$

In this notation  $P = NAT$ . This makes (1) evident.

In the case of a degenerate orbit, we have the following diffeomorphism

$$\mathcal{O}_{\mu_0} = G_{\mu_0} \backslash G \simeq P_{\mu_0} \backslash G^{\mathbb{C}} \tag{2}$$

where  $G_{\mu_0}$  is the stability subgroup and  $P_{\mu_0}$  is the parabolic subgroup with respect to  $\mathcal{O}_{\mu_0}$ . Then  $P_{\mu_0} = NAG_{\mu_0}$ , that proves (2).

On the other hand,  $G$  admits a Gauss decomposition (for the generic type of orbits)

$$G^{\mathbb{C}} = NT^{\mathbb{C}}Z$$

where  $T^{\mathbb{C}}$  is the maximal torus of  $G^{\mathbb{C}}$ , and  $T^{\mathbb{C}} = AT$  in the above notation;  $N$  and  $Z \simeq N^*$  are nilpotent subgroups of  $G^{\mathbb{C}}$  normalized by  $T^{\mathbb{C}}$ . In terms of the root vectors introduced above

$$Z = \exp \left\{ \sum_{\alpha \in \Delta^+} z_\alpha X_{-\alpha} \right\}, \quad z_\alpha \in \mathbb{C}.$$

After [4] we call  $a_\alpha, n_\alpha, z_\alpha$  the **canonical coordinates** connected with the root basis  $\{H_\alpha, X_\alpha, X_{-\alpha}; \alpha \in \Delta^+\}$ . These are coordinates in the group  $G$ .

A comparison of the Gauss and Iwasawa decompositions implies that the orbit  $\mathcal{O}$  is diffeomorphic to the subgroup manifold  $Z$

$$\mathcal{O} \simeq \frac{NAG}{NAT} \simeq \frac{NATZ}{NAT} \simeq Z. \tag{3}$$

Diffeomorphism (3) asserts that one can parameterize the orbit  $\mathcal{O}$  in terms of the complex coordinates  $\{z_\alpha, \alpha \in \Delta^+\}$  that are canonical coordinates in  $Z$ .

However, a Gauss decomposition is local. Therefore, we use a Gauss-Bruhat decomposition instead

$$G^{\mathbb{C}} = \bigcap_{w \in W(G)} PZw.$$

It gives a system of local charts on the orbit

$$\mathcal{O} = P \backslash G^{\mathbb{C}} = \bigcap_{w \in W(G)} Zw. \tag{4}$$

In the case of a degenerate orbit  $\mathcal{O}_{\mu_0}$ ,  $T$  is to be replaced by  $G_{\mu_0}$ , and  $P$  by  $P_{\mu_0}$ . It is sufficient to take the intersection over  $w \in W(G_{\mu_0}) \backslash W(G)$  in (4). Furthermore, in this case,  $Z$  has a less number of coordinates.

**Proposition 3.** *Each orbit  $\mathcal{O}$  of a compact semisimple Lie group  $G$  is locally parameterized in terms of the canonical coordinates  $\{z_\alpha, \alpha \in \Delta^+\}$  in a nilpotent subgroup  $Z$  of  $G^\mathbb{C}$  according to (4).*

Now we apply the above scheme to compact classical Lie groups, namely  $SO(n)$ ,  $SU(n)$ ,  $Sp(n)$ . The scheme consists of several steps. First we parameterize the subgroups  $N$ ,  $A$ , and the group  $G$  in terms of  $\{z_\alpha, \alpha \in \Delta^+\}$ . Secondly, we choose an initial point  $\mu_0$  in the positive closed Weyl chamber  $\overline{C}$  and generate an orbit  $\mathcal{O}_{\mu_0}$  by the dressing formula

$$\hat{\mu} = g^{-1}\mu_0g, \quad g \in G.$$

That gives a parametrization on one of the charts covering the orbit. Finally, we extend the parametrization to all other charts by the action of elements of the Weyl group of  $G$ . We consider the scheme in detail.

**Step 1.** Being a finite group, each classical Lie group has a matrix representation. Let  $\hat{a}$  be the matrix representing an element  $a$ . An Iwasawa decomposition of  $\hat{z} \in Z$  has the following form

$$\hat{z} = \hat{n}\hat{a}\hat{k}, \quad \hat{n} \in N, \quad \hat{a} \in A, \quad \hat{k} \in G. \tag{5}$$

One has to solve (5) in terms of the complex coordinates  $z_\alpha$  that appear as entries of the matrix  $\hat{z}$ . The following transformation of (5) makes the computation easier

$$\hat{z}\hat{z}^* = \hat{n}\hat{a}\hat{k}\hat{k}^*\hat{a}^*\hat{n}^* = \hat{n}\hat{a}^2\hat{n}^*$$

where  $\hat{k}^*$  denotes the hermitian conjugate of  $\hat{k}$ . Indeed,  $\hat{k}\hat{k}^* = e$  for all of the mentioned groups. This is evident, if one considers the conjugation over the complex field in the case of  $SU(n)$ , and over the quaternionic ring in the case of  $Sp(n)$ . If  $\hat{k} \in SO(n)$  one has  $\hat{k}^* = \hat{k}^T$ , and the equality  $\hat{k}\hat{k}^* = e$  is obvious. Moreover, it can easily be checked that  $\hat{a}\hat{a}^* = \hat{a}^2$ . When  $\hat{n}$  and  $\hat{a}$  are parameterized in terms of  $\{z_\alpha\}$ , the matrix  $\hat{k}(z)$  is computed by the formula

$$\hat{k}(z) = \hat{a}^{-1}(z)\hat{n}^{-1}(z)\hat{z}.$$

Here we obtain complex parameterizations of  $N$ ,  $A$ ,  $G$  for all classical compact groups of small dimensions.

**Example 3.** In the case of group  $SU(n)$ , the corresponding complexified group is  $SL(n, \mathbb{C})$ . The subgroup  $N$  consists of complex upper triangular matrices with ones on the diagonal, the subgroup  $Z$  consists of complex low triangular matrices with ones on the diagonal, the subgroup  $A$  contains real diagonal matrices  $\hat{a} = \text{diag}(r_1, r_2, \dots, r_n)$  such that  $\prod_{i=1}^n r_i = 1$ .



Decomposition (5) for a generic orbit  $\mathcal{O}^{\text{SU}(3)}$  gets the form

$$\begin{pmatrix} 1 & 0 & 0 \\ z_1 & 1 & 0 \\ z_3 & z_2 & 1 \end{pmatrix} = \begin{pmatrix} 1 & n_1 & n_3 \\ 0 & 1 & n_2 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \frac{1}{r_1} & 0 & 0 \\ 0 & \frac{r_1}{r_2} & 0 \\ 0 & 0 & r_2 \end{pmatrix} \hat{u}, \quad \hat{u} \in \text{SU}(3)$$

whence it follows

$$r_1^2 = 1 + |z_1|^2 + |z_3 - z_1 z_2|^2, \quad r_2^2 = 1 + |z_2|^2 + |z_3|^2$$

$$n_1 = \frac{1}{r_1^2}(\bar{z}_1(1 + |z_2|^2) - z_2 \bar{z}_3), \quad n_2 = \frac{1}{r_2^2}(\bar{z}_2 + z_1 \bar{z}_3), \quad n_3 = \frac{\bar{z}_3}{r_2^2}.$$

The dressing matrix  $\hat{u}$  is

$$\hat{u} = \begin{pmatrix} \frac{1}{r_1} & -\frac{\bar{z}_1}{r_1} & -\frac{\bar{z}_3 - \bar{z}_1 \bar{z}_2}{r_1} \\ \frac{z_1(1 + |z_2|^2) - z_3 \bar{z}_2}{r_1 r_2} & \frac{1 + |z_3|^2 - z_1 z_2 \bar{z}_3}{r_1 r_2} & -\frac{\bar{z}_2 + z_1 \bar{z}_3}{r_1 r_2} \\ \frac{z_3}{r_2} & \frac{z_2}{r_2} & \frac{1}{r_2} \end{pmatrix}.$$

The case of a degenerate orbit  $\mathcal{O}_d^{\text{SU}(3)}$  is derived from the above by assigning  $z_1 = 0$ , or  $z_2 = 0$ .

**Example 4.** In the case of group  $\text{Sp}(n)$ , the complexified group is  $\text{Sp}(n, \mathbb{C})$ . The both groups describe linear transformations of the quaternionic vector space  $\mathbb{H}^n$ . Therefore, it is suitable to operate with quaternions instead of complex numbers. Each quaternion  $q$  is determined by two complex numbers  $z_1, z_2$  as  $q = z_1 + z_2 \mathbf{j}$ . The quaternionic conjugate of  $q$  is  $\bar{q} = \bar{z}_1 - \mathbf{j} \bar{z}_2$ , where  $\bar{z}_1, \bar{z}_2$  are the complex conjugates of  $z_1, z_2$ . Several useful relations are available

$$\mathbf{j}z = \bar{z}\mathbf{j}, \quad \overline{z + w} = \bar{z} + \bar{w}, \quad \overline{z \cdot w} = \bar{w} \cdot \bar{z}$$

where  $z, w \in \mathbb{C}$ .

The subgroups  $N, Z$  have the same representatives as in the case of group  $\text{SU}(n)$ , but over the quaternionic ring. The subgroup  $A$  consists of real diagonal matrices with the same property as in the case of  $\text{SU}(n)$ .

We start with the simplest group  $\text{Sp}(2)$ . Suppose  $v, q \in \mathbb{H}$  such that  $v = n_1 + n_2 \mathbf{j}$ ,  $q = z_1 + z_2 \mathbf{j}$ , where  $n_1, n_2, z_1, z_2 \in \mathbb{C}$ . Decomposition (5) for an orbit  $\mathcal{O}^{\text{Sp}(2)}$  gets the following form

$$\begin{pmatrix} 1 & 0 \\ q & 1 \end{pmatrix} = \begin{pmatrix} 1 & v \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \frac{1}{r} & 0 \\ 0 & r \end{pmatrix} \hat{p}, \quad \hat{p} \in \text{Sp}(2)$$

whence it follows  $r^2 = 1 + |q|^2$ ,  $v = \bar{q}/r^2$ , or in terms of complex coordinates

$$r^2 = |z_1|^2 + |z_2|^2, \quad n_1 = \frac{\bar{z}_1}{r^2}, \quad n_2 = -\frac{z_2}{r^2}.$$

The dressing matrix  $\hat{p}$  is

$$\hat{p} = \frac{1}{\sqrt{|z_1|^2 + |z_2|^2}} \begin{pmatrix} 1 & -\bar{z}_1 + \mathbf{j}\bar{z}_2 \\ z_1 + z_2\mathbf{j} & 1 \end{pmatrix}.$$

In the case of group  $\text{Sp}(3)$ , we perform all computations in terms of quaternions. Suppose  $q_1 = z_1 + z_2\mathbf{j}$ ,  $q_2 = z_3 + z_4\mathbf{j}$ ,  $q_3 = z_5 + z_6\mathbf{j}$ ,  $v_1 = n_1 + n_2\mathbf{j}$ ,  $v_2 = n_3 + n_4\mathbf{j}$ ,  $v_3 = n_5 + n_6\mathbf{j}$ . Then, for a generic orbit  $\mathcal{O}^{\text{Sp}(3)}$ , one obtains

$$\begin{pmatrix} 1 & 0 & 0 \\ q_1 & 1 & 0 \\ q_3 & q_2 & 1 \end{pmatrix} = \begin{pmatrix} 1 & v_1 & v_3 \\ 0 & 1 & v_2 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \frac{1}{r_1} & 0 & 0 \\ 0 & \frac{r_1}{r_2} & 0 \\ 0 & 0 & r_2 \end{pmatrix} \hat{p}, \quad \hat{p} \in \text{Sp}(3)$$

whence it follows

$$\begin{aligned} r_1^2 &= 1 + |q_1|^2 + |q_3 - q_2q_1|^2, & r_2^2 &= 1 + |q_2|^2 + |q_3|^2 \\ v_1 &= \frac{1}{r_1^2}(\bar{q}_1(1 + |q_2|^2) - \bar{q}_3q_2), & v_2 &= \frac{1}{r_2^2}(\bar{q}_2 + q_1\bar{q}_3), & v_3 &= \frac{\bar{q}_3}{r_2^2}. \end{aligned}$$

The dressing matrix  $\hat{p}$  is

$$\hat{p} = \begin{pmatrix} \frac{1}{r_1} & -\frac{\bar{q}_1}{r_1} & -\frac{\bar{q}_3 - \bar{q}_1\bar{q}_2}{r_1} \\ \frac{q_1(1 + |q_2|^2) - \bar{q}_2q_3}{r_1r_2} & \frac{1 + |q_3|^2 - q_1\bar{q}_3q_2}{r_1r_2} & -\frac{\bar{q}_2 + q_1\bar{q}_3}{r_1r_2} \\ \frac{q_3}{r_2} & \frac{q_2}{r_2} & \frac{1}{r_2} \end{pmatrix}.$$

The case of  $\text{Sp}(n)$  in terms of quaternions is very similar to the case of  $\text{SU}(n)$ . The only warning is that the multiplication of quaternions is not commutative.

**Example 5.** In the case of group  $\text{SO}(n)$ , the corresponding complexified group is  $\text{SO}(n, \mathbb{C})$ . Representatives of the subgroups  $N$  and  $Z$  have not so clear structure as for groups  $\text{SU}(n)$  and  $\text{Sp}(n)$ . The real abelian subgroup  $A$  consists of block-diagonal matrices  $\hat{a} = \text{diag}(A_1, A_2, \dots, A_m)$  in the case of group  $\text{SO}(2m)$ , and  $\hat{a} = \text{diag}(A_1, A_2, \dots, A_m, 1)$  in the case of group  $\text{SO}(2m + 1)$ . Here

$$A_i = \begin{pmatrix} \cosh a_i & -i \sinh a_i \\ i \sinh a_i & \cosh a_i \end{pmatrix}.$$

Consider the group  $\text{SO}(3)$ . The only type of orbits is  $\mathcal{O}^{\text{SO}(3)} = \text{SO}(2) \backslash \text{SO}(3)$ . In this case the decomposition (5) gets the form

$$\begin{pmatrix} 1 - \frac{z^2}{2} & -\frac{iz^2}{2} & -z \\ -\frac{iz^2}{2} & 1 + \frac{z^2}{2} & -iz \\ z & iz & 1 \end{pmatrix} = \begin{pmatrix} 1 - \frac{n^2}{2} & \frac{in^2}{2} & n \\ \frac{in^2}{2} & 1 + \frac{n^2}{2} & -in \\ -n & in & 1 \end{pmatrix} \begin{pmatrix} \cosh a & -i \sinh a & 0 \\ i \sinh a & \cosh a & 0 \\ 0 & 0 & 1 \end{pmatrix} \hat{o}$$

where  $\hat{o} \in \text{SO}(3)$ , and  $a, n, z$  are canonical coordinates in the group. One easily computes the following

$$e^a = 1 + |z|^2, \quad n = \frac{\bar{z}}{1 + |z|^2}.$$

The dressing matrix  $\hat{o}$  is

$$\hat{o} = \begin{pmatrix} \frac{2-z^2-\bar{z}^2}{2(1+|z|^2)} & \frac{i(\bar{z}^2-z^2)}{2(1+|z|^2)} & -\frac{z+\bar{z}}{1+|z|^2} \\ \frac{i(\bar{z}^2-z^2)}{2(1+|z|^2)} & \frac{2+z^2+\bar{z}^2}{2(1+|z|^2)} & -\frac{i(z-z\bar{z})}{1+|z|^2} \\ \frac{z+z\bar{z}}{1+|z|^2} & \frac{i(z-\bar{z})}{1+|z|^2} & \frac{1-|z|^2}{1+|z|^2} \end{pmatrix}.$$

We return to the scheme.

**Step 2.** Suppose we have some parametrization of the dual space  $\mathfrak{g}^*$  of the group  $G$ . We call these parameters group coordinates. In order to parameterize an orbit of  $G$  we find expressions for the group coordinates in terms of the complex coordinates  $\{z_\alpha, \alpha \in \Delta^+\}$ . Let us continue with the example of group  $\text{SU}(3)$ .

Let  $\lambda_a, a = 1, \dots, 8$ , be Gell-Mann matrices, then  $Y_a = -\frac{i}{2}\lambda_a, a = 1, \dots, 8$ , form a basis for  $\mathfrak{g}^*$ . Define a bilinear form on  $\mathfrak{g}^*$  as  $\langle A, B \rangle = -2 \text{Tr } AB$ . Each basis element  $Y_a$  is assigned to a group coordinate:  $\mu_a = \langle \hat{\mu}, Y_a \rangle$ , where

$$\hat{\mu} = -\frac{i}{2} \begin{pmatrix} \mu_3 + \frac{1}{\sqrt{3}}\mu_8 & \mu_1 - i\mu_2 & \mu_4 - i\mu_5 \\ \mu_1 + i\mu_2 & -\mu_3 + \frac{1}{\sqrt{3}}\mu_8 & \mu_6 - i\mu_7 \\ \mu_4 + i\mu_5 & \mu_6 + i\mu_7 & -\frac{2}{\sqrt{3}}\mu_8 \end{pmatrix}.$$

A coadjoint orbit is generated by the dressing formula

$$\hat{\mu} = \hat{u}^* \hat{\mu}_0 \hat{u}, \quad \hat{\mu}_0 \in \mathfrak{h}^*$$

where  $\hat{\mu}_0$  is an initial point. As shown in Section 2, each orbit is uniquely defined by a point of the closed positive Weyl chamber. Let simple roots of  $\mathfrak{su}(3)$  be as follows:  $\hat{\alpha}_1 = \text{diag}(i, -i, 0)$  and  $\hat{\alpha}_2 = \text{diag}(0, i, -i)$ . The closed positive Weyl chamber is the set of points  $\hat{\mu}_0$  such that

$$\hat{\mu}_0 = -\frac{i}{3}\xi \begin{pmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix} - \frac{i}{3}\eta \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix}, \quad \xi, \eta > 0. \quad (6)$$

Obviously, walls of the Weyl chamber are obtained by assigning  $\xi = 0$  or  $\eta = 0$ . In this notation  $\Gamma_{\alpha_1} = \{-\frac{i}{3}\eta \text{diag}(1, 1, -2); \eta > 0\}, \Gamma_{\alpha_2} = \{-\frac{i}{3}\xi \text{diag}(2, -1, -1); \xi > 0\}$ . The chosen representation of an initial point  $\hat{\mu}_0$  is the most suitable for the further computation.

According to Proposition 1 we get a generic orbit if  $\eta \neq 0$  and  $\xi \neq 0$ . If  $\xi$  or  $\eta$  vanishes, we get a degenerate one. A generic orbit is parameterized by three

complex coordinates  $z_1, z_2, z_3$ . If  $\xi$  vanishes, one has to assign  $z_1 = 0$ . If  $\eta$  vanishes, then  $z_2 = 0$ . We consider the degenerate orbit through the following point

$$\hat{\mu}_0 = -\frac{i}{3}\eta \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix}.$$

One can attach some physical meaning to nonzero entries of the initial point  $\hat{\mu}_0$  because of its diagonal form. For in quantum mechanics diagonal matrices represent observable variables. Suppose  $\hat{\mu}_0$  is the value of  $\hat{\mu}$  at the infinity:  $\hat{\mu}_0 = \hat{\mu}(\infty)$ . The diagonal entries are expressed in terms of the group coordinates  $\mu_3$  and  $\mu_8$  and we fix their values at the infinity as follows:  $\mu_3(\infty) = m, \mu_8(\infty) = q$ . Then

$$\eta = -\frac{1}{2}(m - \sqrt{3}q), \quad \xi = m.$$

Suppose the group  $\text{SU}(3)$  describes a magnetic with spin one. Then  $m$  serves as a projection of magnetic moment (magnetization) of the magnetic, and  $q$  serves as a projection of quadrupole moment.

The dressing procedure gives the following explicit expression for the generalized stereographic projection onto a generic orbit of  $\text{SU}(3)$

$$\begin{aligned} \mu_1 &= -\frac{\eta}{r_2^2}(\bar{z}_2 z_3 + z_2 \bar{z}_3) - \frac{\xi}{r_1^2}(z_1 + \bar{z}_1) \\ \mu_2 &= \frac{i\eta}{r_2^2}(\bar{z}_2 z_3 - z_2 \bar{z}_3) + \frac{i\xi}{r_1^2}(z_1 - \bar{z}_1) \\ \mu_3 &= \frac{\eta}{r_2^2}(|z_2|^2 - |z_3|^2) + \frac{\xi}{r_1^2}(1 - |z_1|^2) \\ \mu_4 &= -\frac{\eta}{r_2^2}(z_3 + \bar{z}_3) - \frac{\xi}{r_1^2}(z_3 - z_1 z_2 + \bar{z}_3 - \bar{z}_1 \bar{z}_2) \\ \mu_5 &= \frac{i\eta}{r_2^2}(z_3 - \bar{z}_3) + \frac{i\xi}{r_1^2}(z_3 - z_1 z_2 - (\bar{z}_3 - \bar{z}_1 \bar{z}_2)) \\ \mu_6 &= -\frac{\eta}{r_2^2}(z_2 + \bar{z}_2) + \frac{\xi}{r_1^2}(\bar{z}_1(z_3 - z_1 z_2) + z_1(\bar{z}_3 - \bar{z}_1 \bar{z}_2)) \\ \mu_7 &= \frac{i\eta}{r_2^2}(z_2 - \bar{z}_2) - \frac{i\xi}{r_1^2}(\bar{z}_1(z_3 - z_1 z_2) - z_1(\bar{z}_3 - \bar{z}_1 \bar{z}_2)) \\ \sqrt{3}\mu_8 &= \frac{\eta}{r_2^2}(2 - |z_2|^2 - |z_3|^2) + \frac{\xi}{r_1^2}(1 + |z_1|^2 - 2|z_3 - z_1 z_2|^2) \end{aligned} \tag{7}$$

where

$$r_1^2 = 1 + |z_1|^2 + |z_3 - z_1 z_2|^2, \quad r_2^2 = 1 + |z_2|^2 + |z_3|^2.$$

Obviously, all expressions can be divided into two parts: with the coefficients  $\eta$  and  $\xi$ . These parts correspond to the basis matrices in (6).

For the stereographic projection onto a degenerate orbit through  $\hat{\mu}_0$  chosen above one has to assign  $\xi = 0, z_1 = 0$  in (7).

**Step 3.** Parametrization (7) is available on the coordinate chart containing the point  $(z_1 = 0, z_2 = 0, z_3 = 0)$ . By the action of elements of the Weyl group one obtains parameterizations on all other charts. The Weyl group is generated by reflections across the hyperplanes orthogonal to simple roots. In the case of group  $SU(3)$ , these reflections are represented by the following matrices

$$\hat{w}_1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad \hat{w}_2 = \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}.$$

The action of  $\hat{w}_1$  transforms the chart with coordinates (7) onto another one by the following change of coordinates

$$(z_1, z_2, z_3) \mapsto (z'_1, z'_2, z'_3), \quad z'_1 = \frac{1}{z_1}, \quad z'_2 = -z_3, \quad z'_3 = -z_2.$$

This chart contains the point  $(z_1 = \infty, z_2 = 0, z_3 = 0)$ . The action of  $\hat{w}_2$  transforms coordinates (7) by the following change of coordinates

$$(z_1, z_2, z_3) \mapsto (z'_1, z'_2, z'_3), \quad z'_1 = -(z_3 - z_1 z_2), \quad z'_2 = \frac{1}{z_2}, \quad z'_3 = -\frac{z_3}{z_2}.$$

The latter chart contains the point  $(z_1 = 0, z_2 = \infty, z_3 = 0)$ .

Evidently, the other elements of  $W(SU(3))$  are  $\hat{e}, \hat{w}_1 \hat{w}_2, \hat{w}_2 \hat{w}_1, \hat{w}_1 \hat{w}_2 \hat{w}_1$ . The corresponding changes of coordinates are obtained by sequential actions of the two described above.

#### 4. Kählerian Structure on the Coadjoint Orbits

The perfect property of coadjoint orbits of compact semisimple Lie groups is the following. Each orbit is simultaneously a Riemannian manifold and a symplectic one. A Riemannian metrics and the matched symplectic form together are called a **Kählerian structure**. Borel [5] proved the following

**Proposition 4.** *Suppose  $G$  is a semisimple compact Lie group. Then each orbit of  $G$  admits a complex analytic Kählerian structure invariant under the group  $G$ .*

It means that each orbit possesses a hermitian Riemannian metrics, the Kählerian metrics  $ds^2$ , and the corresponding closed two-form, the **Kählerian form**  $\omega$

$$ds^2 = \sum_{\alpha, \beta} g_{\alpha\bar{\beta}} dz_\alpha d\bar{z}_\beta, \quad \omega = \sum_{\alpha, \beta} ig_{\alpha\bar{\beta}} dz_\alpha \wedge d\bar{z}_\beta.$$

The  $G$ -invariance of a Kählerian structure means invariance under the action of  $G$ . Here we consider the action of a group as right multiplication. A Kählerian

structure is determined by a **Kählerian potential**  $\Phi$  according to the formula

$$g_{\alpha\bar{\beta}} = \frac{\partial^2 \Phi}{\partial z_\alpha \partial \bar{z}_\beta}, \quad \omega_{\alpha\bar{\beta}} = ig_{\alpha\bar{\beta}}.$$

The objective of this section is to obtain an expression for a Kählerian structure on a coadjoint orbit. Evidently, for this purpose it is sufficient to find a Kählerian potential, which simultaneously gives the Kählerian metrics and the Kählerian form.

On the other hand, one has the following

**Proposition 5** (see [6]). *If  $G$  is a compact semisimple Lie group, the Kirillov-Kostant-Souriau two-form coincides with a  $G$ -invariant Kählerian form.*

While we deal with compact semisimple classical Lie groups, we can use a Kirillov-Kostant-Souriau differential form as a Kählerian form.

Define a bilinear form on  $\mathfrak{g}$  as follows

$$\langle X, Y \rangle = \text{Tr } XY, \quad X, Y \in \mathfrak{g}.$$

In the case of classical Lie groups, the bilinear form is proportional to the standard Killing form on  $\mathfrak{g}$ .

Define a vector field  $\tilde{X}$  on a coadjoint orbit  $\mathcal{O}$  by

$$\tilde{X}f(\mu) = \frac{d}{d\tau} f(\text{Ad}_{\exp(\tau X)}^* \mu) \Big|_{\tau=0}, \quad f \in C^\infty(\mathcal{O}).$$

One can introduce an Ad-invariant closed two-form on  $\mathcal{O}$  by the formula

$$\omega(\tilde{X}, \tilde{Y}) = \langle \mu, [X, Y] \rangle, \quad X, Y \in \mathfrak{g}, \quad \mu \in \mathfrak{g}^*. \tag{8}$$

This two-form is called a **Kirillov-Kostant-Souriau form**.

The straightforward way of obtaining a Kählerian form is to solve equations (8). Unfortunately, it becomes extremely complicate in dimensions greater than three. This way is developed by Picken [3]. He computes Kählerian forms on flag manifolds via  $G$ -invariant one-forms in terms of Bruhat coordinates.

We return to the idea of finding a Kählerian potential instead of a Kählerian form. In general, each  $G$ -covariant real function on an orbit serves as a Kählerian potential. It turns out, that each orbit has a unique  $G$ -covariant real function, which we call a **Kählerian potential on the orbit**.

The same idea is used by Alekseevsky and Perelomov [7]. In order to find potentials for all closed two-forms on orbits of group  $GL(n)$ , they consider the real positive functions built by means of principal minors of  $\hat{z}\hat{z}^* \in GL(n)$ , and select the functions that are  $G$ -covariant. Here we develop the idea of Alekseevsky and Perelomov, because this way allows to avoid complicate computations.

Below we prove that a Kählerian potential is determined by an one-dimensional irreducible representation of the real abelian subgroup  $A$  of  $G^{\mathbb{C}}$ . We use the group-theoretical approach in our proof.

Each orbit  $\mathcal{O} = P \backslash G^{\mathbb{C}}$  is a holomorphic manifold, which admits the construction of a line bundle. Let  $\{U_k\}$  be its atlas. An arbitrary  $g_{\mathbb{C}} \in G^{\mathbb{C}}$  has a decomposition

$$g_{\mathbb{C}} = h_k(x)s_k(x), \quad x \in U_k \tag{9}$$

where  $s_k : U_k \rightarrow G^{\mathbb{C}}$  is a local section of  $\mathcal{O}$ . If  $U_k \cap U_j \neq \emptyset$ , then there exists a map  $s_{kj} = s_k \circ s_j^{-1}$ , which is  $s_{kj} : U_k \cap U_j \rightarrow P$ . A one-dimensional representation of the parabolic subgroup  $P$  of  $G^{\mathbb{C}}$  gives a  $G$ -covariant function on an orbit.

Recall, that  $P = NAT$  in the case of a generic orbit. In the case of a degenerate orbit, one has  $P = NAG_{\mu_0}$ , where  $G_{\mu_0}$  is the stability subgroup at an initial point  $\mu_0 \in \mathfrak{h}^*$  giving rise to the orbit. A one-dimensional irreducible representation is trivial on any nilpotent group. This means that the representation of  $P$  coincides with the representation of the maximal torus  $T^{\mathbb{C}} = AT$  of  $G^{\mathbb{C}}$ . Moreover, we are interested in real representations because a Kählerian potential is a real function. Consequently, the required representation is determined only by  $A$ .

Now we build a one-dimensional irreducible representation of  $T^{\mathbb{C}}$ . Obviously,  $T^{\mathbb{C}}$  is isomorphic to a direct product of  $l$  samples of the multiplicative group  $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$ , where  $l = \dim T$ . Let the following set of complex numbers  $(d_1, d_2, \dots, d_l)$  be an image of  $\hat{d} \in T^{\mathbb{C}}$  under the isomorphism. It is clear that the set of real numbers  $(r_1, r_2, \dots, r_l)$ , where  $r_i = |d_i|$ ,  $i = 1, \dots, l$ , is an image of  $\hat{a} \in A$  under the isomorphism. In terms of complex coordinates  $z = \{z_{\alpha}; \alpha \in \Delta^+\}$ , which are canonical coordinates in  $Z$ , an Iwasawa decomposition of any  $\hat{z} \in Z$  gets the form

$$\hat{z} = \hat{n}(z)\hat{a}(z)\hat{k}(z). \tag{10}$$

Here  $\hat{k}(z)$  represents a point of the orbit in terms of the complex coordinates  $\{z_{\alpha}\}$  while  $\hat{n}(z)$  and  $\hat{a}(z)$  denote matrices  $\hat{n}$  and  $\hat{a}$  in terms of  $\{z_{\alpha}\}$ . After the action of an element  $g \in G$  on  $z$  we perform a Gauss-Bruhat decomposition

$$\hat{z}\hat{g} = \hat{n}_B(zg)\hat{d}(zg)\hat{z}_g, \quad \hat{n}_B(zg) \in N. \tag{11}$$

From the Iwasawa decomposition of  $\hat{z}_g$  we have

$$\hat{a}(z_g) = \hat{n}^{-1}(z_g)\hat{z}_g\hat{k}^{-1}(z_g).$$

Using (10) and (11) we get

$$\hat{a}(z_g) = \hat{n}^{-1}(z_g)\hat{d}^{-1}(zg)\hat{n}_B^{-1}(zg)\hat{n}(z)\hat{a}(z)\hat{k}(z)\hat{g}\hat{k}^{-1}(z_g). \tag{12}$$

In order to gather nilpotent elements together we recall that the maximal torus  $T^{\mathbb{C}}$  is the normalizer of  $N$ , that gives the following equality

$$\hat{d}^{-1}(zg)\hat{n}_B^{-1}(zg)\hat{n}(z)\hat{d}(zg) = \hat{n}(z, g), \quad \hat{n}(z, g) \in N.$$

Substituting  $\hat{n}_B(zg)\hat{d}^{-1}(zg)$  for  $\hat{d}^{-1}(zg)\hat{n}_B^{-1}(zg)\hat{n}(z)$  in (12) we obtain

$$\hat{a}(z_g) = \hat{n}^{-1}(z_g)\hat{n}(z, g)\hat{d}^{-1}(zg)\hat{a}(z)\hat{k}(z)\hat{g}\hat{k}^{-1}(z_g).$$

To cancel the element  $\hat{k}(z)\hat{g}\hat{k}^{-1}(z_g) = \hat{g}' \in G$  we take the following product

$$\hat{a}^2(z_g) = \hat{a}(z_g)\hat{a}^*(z_g) = \hat{n}\hat{d}^{-1}(z_g)\hat{a}^2(z)\hat{d}^{*-1}(z_g)\hat{n}^* \tag{13}$$

where  $\hat{n}$  denotes  $\hat{n}^{-1}(z_g)\hat{n}(z, g) \in N$ .

Now we construct a one-dimensional real representation of (13). Let  $\chi^\xi(\hat{a})$  denote a representation of  $\hat{a}$  with real weights  $\xi = (\xi_1, \xi_2, \dots, \xi_l)$ . A one-dimensional real representation of  $\hat{a} \in A$  has the following form  $\chi^\xi(\hat{a}) = r_1^{\xi_1} r_2^{\xi_2} \dots r_l^{\xi_l}$ , and a one-dimensional real representation of  $\hat{d} \in T^C$  has the form  $\chi^\xi(\hat{d}) = d_1^{\xi_1} d_2^{\xi_2} \dots d_l^{\xi_l}$ . Therefore, the representation of  $\hat{a}^2(z_g)$  gets the form

$$\chi^{2\xi}(\hat{a}(z_g)) = \chi^\xi(\hat{d}(zg)) \overline{\chi^\xi(\hat{d}(zg))} \chi^{2\xi}(\hat{a}(z)).$$

Whence it is seen that  $\chi^{2\xi}(\hat{a}(z))$  is transformed by a cocycle  $\chi^\xi(\hat{d}(zg))$  defined on  $G \times \mathcal{O}$ . It means that the function

$$\ln \chi^{2\xi}(\hat{a}(z)) = \xi_1 \ln r_1^2(z) + \xi_2 \ln r_2^2(z) + \dots + \xi_l \ln r_l^2(z) \tag{14}$$

is  $G$ -covariant, and serves as a Kählerian potential on  $\mathcal{O}$ . Moreover, each function  $\ln r_i^2(z)$ ,  $i = 1, \dots, l$ , is a Kählerian potential itself.

Remarkably, that each coadjoint orbit has a unique Kählerian potential of the form (14), where the weights  $\xi = (\xi_1, \xi_2, \dots, \xi_l)$  are determined by an initial point of the orbit. We have proven the following

**Proposition 6.** *Suppose  $A$  is the real abelian subgroup of  $G^C$ ,  $\hat{a} \in A$ , and  $\chi^\xi(\hat{a})$  is the one-dimensional representation of  $\hat{a}$  with real weights  $\xi = (\xi_1, \xi_2, \dots, \xi_l)$ . Then Kählerian potentials on coadjoint orbits of  $G$  have the form  $\ln \chi^{2\xi}(\hat{a})$ , moreover each orbit has the Kählerian potential with a unique  $\xi$ .*

**Remark 1.** *In the case of integer weights  $\xi = (\xi_1, \xi_2, \dots, \xi_l)$ , the line bundle over each coadjoint orbit of  $G$  is holomorphic. This idea is derived from the Borel-Weyl theory based on [8].*

Let us consider some examples.

**Example 6.** In the case of group  $SU(n)$ , a representative of the real abelian subgroup  $A$  has the form of a diagonal matrix with  $\det \hat{a} = 1$ , that is

$$\hat{a} = \text{diag}(1/r_1, r_1/r_2, \dots, r_{n-2}/r_{n-1}, r_{n-1})$$

and  $\dim T = n - 1$ . Let  $(r_1, r_2, \dots, r_{n-1})$  be an image of  $\hat{a}$  under an isomorphism from  $T^C$  onto  $(C^*)^{n-1}$ . Then  $\chi^\xi(\hat{a}) = r_1^{\xi_1} r_2^{\xi_2} \dots r_{n-1}^{\xi_{n-1}}$ , where  $\xi_i \in \mathbb{R}$ ,



$i = 1, \dots, n - 1$ , whence Kählerian potentials have the following form

$$\Phi = \xi_1 \ln r_1^2 + \xi_2 \ln r_2^2 + \dots + \xi_{n-1} \ln r_{n-1}^2.$$

For instance, Kählerian potentials on orbits of  $SU(3)$  are

$$\Phi = \xi \ln r_1^2 + \eta \ln r_2^2, \quad r_1^2 = 1 + |z_1|^2 + |z_3 - z_1 z_2|^2, \quad r_2^2 = 1 + |z_2|^2 + |z_3|^2.$$

This expression completely accords with the straightforward solution of (8), which gives the following

$$\begin{aligned} \Phi &= \langle \hat{\mu}_0, \hat{\alpha}_1 \rangle \Phi_1 + \langle \hat{\mu}_0, \hat{\alpha}_2 \rangle \Phi_2 \\ \Phi_1 &= \ln(1 + |z_1|^2 + |z_3 - z_1 z_2|^2), \quad \Phi_2 = \ln(1 + |z_2|^2 + |z_3|^2). \end{aligned}$$

Here  $\hat{\mu}_0$  is an initial point of an orbit and  $\hat{\alpha}_1, \hat{\alpha}_2$  are the simple roots of  $\mathfrak{su}(3)$ . In the case of a degenerate orbit, one has to assign  $z_1 = 0$  or  $z_2 = 0$ .

**Example 7.** In the case of groups  $SO(n)$ ,  $n = 2m$  and  $n = 2m + 1$ , a representative of the subgroup  $A$  has the form of a block-diagonal matrix, namely

$$\hat{a} = \text{diag}(A_1, A_2, \dots, A_m) \quad \text{or} \quad \hat{a} = \text{diag}(A_1, A_2, \dots, A_m, 1)$$

$$A_i = \begin{pmatrix} \cosh a_i & -i \sinh a_i \\ i \sinh a_i & \cosh a_i \end{pmatrix}, \quad i = 1, \dots, m.$$

Here  $\{a_i\}$  are canonical coordinates in the maximal torus  $T$ , and  $\dim T = m$ . Let  $(e^{a_1}, e^{a_2}, \dots, e^{a_m})$  be an image of  $\hat{a}$  under an isomorphism from  $T^{\mathbb{C}}$  onto  $(\mathbb{C}^*)^m$ . Then  $\chi^\xi(\hat{a}) = e^{\xi_1 a_1} e^{\xi_2 a_2} \dots e^{\xi_m a_m}$ , whence it follows

$$\Phi = 2\xi_1 a_1 + 2\xi_2 a_2 + \dots + 2\xi_m a_m.$$

Kählerian potentials on coadjoint orbits of  $SO(4)$  computed by (8) have the form

$$\begin{aligned} \Phi &= \langle \hat{\mu}_0, \hat{\alpha}_1 \rangle \Phi_1 + \langle \hat{\mu}_0, \hat{\alpha}_2 \rangle \Phi_2 \\ \Phi_1 &= \ln(1 + |z_1|^2) - \ln(1 + |z_2|^2), \quad \Phi_2 = \ln(1 + |z_1|^2) + \ln(1 + |z_2|^2). \end{aligned}$$

Here the bilinear form on  $\mathfrak{so}(4)$  is defined by  $\langle A, B \rangle = \frac{1}{2} \text{Tr } AB$ .

**Proposition 7.** *The Kählerian potential on each coadjoint orbit  $\mathcal{O}_{\mu_0}$  of a compact classical Lie group  $G$  has the following form*

$$\Phi = \sum_k \langle \mu_0, \alpha_k \rangle \Phi_k, \quad \Phi_k = a_{\alpha_k}$$

where  $\alpha_k$  is a simple root of  $\mathfrak{g}$ ,  $a_{\alpha_k}$  is the canonical coordinate corresponding to  $H_{\alpha_k} \in \mathfrak{h}$ , and  $\langle \cdot, \cdot \rangle$  denotes a bilinear form on the dual space of  $\mathfrak{g}$ .

**Remark 2.** *If  $\mu_0$  satisfies the integer condition*

$$2 \frac{\langle \mu_0, \alpha_k \rangle}{\langle \alpha_k, \alpha_k \rangle} \in \mathbb{Z}$$

for all simple roots  $\alpha_k$  of  $\mathfrak{g}$  then the orbit through  $\mu_0$  can be quantized. In other words, there exists an irreducible unitary representation of  $G$  in the space of holomorphic sections on the orbit. Each section serves as a quantum state.

### 5. Cohomology Rings of Coadjoint Orbits

In the last section we examine the cohomology rings of coadjoint orbits of compact semisimple Lie groups. In [9] Borel proved that all forms of odd degrees on the orbit are precise. Therefore, we are interested in the forms of even degrees. In order to introduce a basis for the cohomology ring it is sufficient to find a basis for the cohomology group  $H^2$ .

In the case of a generic coadjoint orbit of a compact semisimple Lie group  $G$ , the following formula is available

$$b^0 + b^2 + \dots + b^{2n} = \text{ord } W(G)$$

where  $b^k$  denotes the Betti number of a cohomology group  $H^k$ . In the case of a degenerate orbit, one has to modify the formula as

$$b^0 + b^2 + \dots + b^{2m} = \frac{\text{ord } W(G)}{\text{ord } W(G_{\mu_0})}$$

where  $G_{\mu_0}$  is the stability subgroup at  $\mu_0$ .

**Example 8.** In the case of group  $SU(2)$ , we have the only type of orbits  $\mathcal{O}^{SU(2)}$  of dimension two. The Weyl group  $W(SU(2))$  also has dimension two. Therefore, the cohomology ring consists of two cohomology groups, each of dimension one

$$H^* = H^0 \oplus H^2, \quad 1 + 1 = 2.$$

In the case of group  $SU(3)$ , we have two types of orbits: a generic one  $\mathcal{O}^{SU(3)}$  of dimension six, and a degenerate one  $\mathcal{O}_d^{SU(3)}$  of dimension four. In the case of a generic orbit, the Weyl group has dimension six, and the cohomology ring is

$$H^* = H^0 \oplus H^2 \oplus H^4 \oplus H^6, \quad 1 + 2 + 2 + 1 = 6.$$

For a degenerate orbit we have  $\frac{\text{ord } W(G)}{\text{ord } W(G_{\mu_0})} = 3$ , and the cohomology ring is

$$H^* = H^0 \oplus H^2 \oplus H^4, \quad 1 + 1 + 1 = 3.$$

Recall the well-known Leray-Hirsch theorem.

**Theorem (Leray-Hirsch).** *Suppose  $\mathcal{E}$  is a fibre bundle over a base  $\mathcal{M}$  with a fibre  $\mathcal{F}$ , and  $\omega_1, \omega_2, \dots, \omega_r$  are cohomology classes on  $\mathcal{E}$  that being restricted to each fibre give its cohomologies. Then*

$$H^*(\mathcal{E}) = H^*(\mathcal{M}) \otimes H^*(\mathcal{F}).$$

Apply the theorem to an orbit  $\mathcal{O}$  regarded as a fibre bundle over an orbit  $\mathcal{O}_1$  with an orbit  $\mathcal{O}_2$  as a fibre, that is  $\mathcal{O} = \mathcal{E}(\mathcal{O}_1, \mathcal{O}_2, \pi)$ . The cohomology ring of  $\mathcal{O}$  is a tensor product of the cohomology rings of the base and the fiber

$$H * (\mathcal{O}) = H * (\mathcal{O}_1) \otimes H * (\mathcal{O}_2).$$

Conversely, if one finds coherent cohomology classes on  $\mathcal{O}_1$  and  $\mathcal{O}_2$ , then one can construct the cohomology ring of  $\mathcal{O}$  by the latter formula. It means, the cohomology ring of a generic orbit can be derived from the cohomology rings of a degenerate orbit and a generic orbit of a group of less dimension.

**Example 9.** We continue to deal with the group  $SU(3)$ . It was shown that

$$\mathcal{O}^{SU(3)} = \mathcal{E} \left( \mathcal{O}_d^{SU(3)}, \mathcal{O}^{SU(2)}, \pi \right).$$

Then the cohomology ring of  $\mathcal{O}^{SU(3)}$  is the tensor product of the cohomology rings of the orbits  $\mathcal{O}_d^{SU(3)}$  and  $\mathcal{O}^{SU(2)}$

$$\begin{aligned} H^*(\mathcal{O}^{SU(3)}) &= (H^0 \oplus H^2 \oplus H^4) \otimes (H^0 \oplus H^2) \\ &= H^0 \otimes H^0 \oplus \underbrace{H^0 \otimes H^2 \oplus H^2 \otimes H^0}_{H^2(\mathcal{O}^{SU(3)})} \oplus \underbrace{H^2 \otimes H^2 \oplus H^4 \otimes H^0}_{H^4(\mathcal{O}^{SU(3)})} \oplus H^4 \otimes H^2. \end{aligned}$$

Obviously, the cohomology groups  $H^2$  and  $H^4$  of  $\mathcal{O}^{SU(3)}$  both have dimension two. Moreover, from the previous expression we can see the structure of a basis for  $H^2$

$$H^2 \left( \mathcal{O}^{SU(3)} \right) = H^0(1) \otimes H^2(2) \oplus H^2(1) \otimes H^0(2)$$

where 1 denotes  $\mathcal{O}_d^{SU(3)} \simeq \mathbb{C}\mathbb{P}^2$ , and 2 denotes  $\mathcal{O}^{SU(2)} \simeq \mathbb{C}\mathbb{P}^1$ .

At the same time, a suitable basis for  $H^2$  can be obtained from Kählerian potentials on coadjoint orbits of a group. As shown in the previous section, all two-forms on the orbits of a compact classical Lie group  $G$  have the form

$$\omega = \sum_k ic_k \sum_{\alpha, \beta} \frac{\partial^2 \Phi_k}{\partial z_\alpha \partial \bar{z}_\beta} dz_\alpha \wedge d\bar{z}_\beta, \quad k = 1, \dots, \dim T$$

where  $\Phi_k$  coincides with the canonical coordinate  $a_{\alpha_k}$  corresponding to  $H_{\alpha_k} \in \mathfrak{h}$ . Obviously,  $\dim H^2 = \dim T = l$ . Consequently, one can find precisely  $l$  two-forms that give a basis for  $H^2$ .

The standard way to generate a basis for  $H^2$  is the following. Let  $H_2$  be the homology group adjoint to  $H^2$ . By  $[\gamma]$  we denote a class of two-cycles, which can be represented as spheres. The sphere is an orbit of a subgroup  $SU_\alpha(2)$

$$SU_\alpha(2) \simeq \exp\{H_\alpha, (X_\alpha - X_{-\alpha}), i(X_\alpha + X_{-\alpha})\}, \quad \alpha \in \Delta_+.$$

Suppose we find  $l$  independent two-cycles connected with the simple roots of  $\mathfrak{g}$ , we denote them by  $\gamma_i$ . The basis for  $H^2$  consists of two-forms  $\omega_j$  such that

$$\int_{\gamma_i} \omega_j = \delta_{ij} \tag{15}$$

where  $\delta_{ij}$  is the Kronecker symbol.

**Example 10.** We consider coadjoint orbits of  $SU(3)$  as an example. Let the simple roots of  $\mathfrak{su}(3)$  be as follows:  $\alpha_1 = \text{diag}(i, -i, 0)$  and  $\alpha_2 = \text{diag}(0, i, -i)$ . Then the independent two-cycles are generated by the following dressing matrices

$$\hat{u}_1 = \begin{pmatrix} \frac{1}{\sqrt{1+|z_1|^2}} & \frac{-\bar{z}_1}{\sqrt{1+|z_1|^2}} & 0 \\ \frac{z_1}{\sqrt{1+|z_1|^2}} & \frac{1}{\sqrt{1+|z_1|^2}} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \hat{u}_2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{\sqrt{1+|z_2|^2}} & \frac{-\bar{z}_2}{\sqrt{1+|z_2|^2}} \\ 0 & \frac{z_2}{\sqrt{1+|z_2|^2}} & \frac{1}{\sqrt{1+|z_2|^2}} \end{pmatrix}$$

which are obtained from the dressing matrix  $\hat{u}$  by assigning  $z_2 = z_3 = 0$  or  $z_1 = z_3 = 0$ , respectively. The two-forms  $\omega_j$  satisfying (15) are

$$\omega_j = \frac{1}{2\pi} \sum_{\alpha,\beta} \frac{\partial^2 \Phi_j}{\partial z_\alpha \partial \bar{z}_\beta} dz_\alpha \wedge d\bar{z}_\beta, \quad j = 1, 2$$

$$\Phi_1 = \ln(1 + |z_1|^2 + |z_3 - z_1 z_2|^2), \quad \Phi_2 = \ln(1 + |z_2|^2 + |z_3|^2).$$

They form a basis for  $H^2(\mathcal{O}^{SU(3)})$ .

### 6. Conclusion

In this paper we develop a unified approach to solutions of the announced problems for a coadjoint orbit of a compact semisimple classical Lie group  $G$ . The problems are the following: an explicit parametrization of the orbit, obtaining a Kählerian structure, introducing basis forms for the cohomology group of the orbit. The key role belongs to the subgroup  $A$  in an Iwasawa decomposition, this is the real abelian subgroup of a complexification of the group  $G$ . The subgroup  $A$  determines a Kählerian potential on each orbit and a suitable basis for the cohomology group  $H^2$  of the orbit.

Our investigation concerns classical (matrix) Lie groups. The same problems in the general case remain of current importance.

### Acknowledgements

We would like to thank professor Andreas Arvanitoyergos for fruitful debates and proper references, Gadano Vilasi for valuable discussions, and Ivailo Mladenov and Mariana Hadzhilazova for their hospitality.

## References

- [1] Bott R., *The Geometry and Representation Theory of Compact Lie Groups*, In: Representation Theory of Lie Groups, London Mathematical Society Lecture Note Series, vol. 34, Cambridge Univ. Press, Cambridge, 1979, pp 65–90.
- [2] Skrypnyk T., *Coadjoint Orbits of Compact Lie Groups and Generalized Stereographic Projection*, Ukr. Math. J. **51** (1999) 1714–1718.
- [3] Picken R. *The Duistermaat-Heckman Integration Formula on Flag Manifolds*, J. Math. Phys. **31** (1990) 616–638.
- [4] Helgason S., *Differential Geometry, Lie Groups and Symmetric Spaces*, Academic Press, Toronto, 1978.
- [5] Borel A., *Kählerian Coset Spaces of Semisimple Lie Groups*, PNAS **40** (1954) 1147–1151.
- [6] Besse A., *Einstein Manifolds*, Springer, 1987.
- [7] Alekseevsky D. and Perelomov A. *Invariant Kähler-Einstein Metrics on Compact Hologenious Spaces*, Functional Anal. Appl. **20** (1986) 171–182.
- [8] Borel A. and Weyl A., *Representations lineares et espaces homogenes Kählerians des groupes de Lie compacts*, Seminaire Bourbaki, May 1954.
- [9] Borel A., *Sur la cohomologie des espaces fibres principaux et des espaces homogenes de groupes de Lie compacts*, Ann. Math. **57** (1953) 115–207.